

## DYSRHYTHMIA MONITORING

### Expected Practice:

- ☑ Select the best monitoring leads for dysrhythmia identification (display two leads when possible).
  - Lead V<sub>1</sub> to diagnose wide QRS complex.
  - Lead II to diagnose atrial activity and measure heart rate.
- ☑ Proper electrode placement is required for accurate diagnosis (Figure 1).
- ☑ Prepare the patient's skin before attaching ECG electrodes.
- ☑ Measure QT interval and calculate QTc using a consistent lead if high risk for Torsades de Pointes.

### Scope and Impact of the Problem:

- Studies show that nurses often monitor in a single lead regardless of diagnosis.<sup>1-2</sup>
- Failure to properly prep skin prior to electrode placement may cause inappropriate monitoring alarms.<sup>3-4</sup>
- When an electrode is misplaced by as little as 1 intercostal space, QRS morphology can change and misdiagnosis may occur (i.e., ventricular tachycardia [VT] may be misidentified as supraventricular tachycardia [SVT] or vice versa).<sup>5</sup>

### Supporting Evidence:

- V<sub>1</sub> is the lead of choice to diagnose wide QRS complexes (VT vs. SVT with aberrant conduction; left vs. right BBB). A 5 lead monitoring system is required to monitor V leads. MCL<sub>1</sub> may differ in QRS morphology as compared to V<sub>1</sub> and should be used only when a 5 lead monitoring system is unavailable.<sup>6-10</sup> (Level V)
- When V<sub>1</sub> electrode placement is not possible, V<sub>6</sub> may be used.<sup>7,11</sup> (Level IV)
- Electrode site preparation includes clipping excessive hair and cleansing oily skin with alcohol.<sup>3-4</sup> (Level IV)
- QTc >0.50 sec (500 ms) is dangerously prolonged and associated with risk for Torsades de Pointes. The QT interval should be corrected for heart rate (QTc) and monitored with any of the following:<sup>9-10,12-15</sup> (Level IV)
  - Antidysrhythmic, antibiotic, antipsychotic, and other drugs that prolong QTc
  - Severe bradycardia
  - Hypokalemia or hypomagnesemia
  - Any drug overdose
- Perform an atrial electrogram (AEG) in cardiac surgical patients with atrial epicardial wires to assist in identifying atrial activity.<sup>16-17</sup> (Level V)




Figure 1

### **Pediatric Specific**

Abnormal prolongation: QTc >0.40 sec ± 10%. Pediatric limits are age specific and shorter than adult ranges.<sup>18</sup>

### Actions for Nursing Practice:

- Ensure that your organization has written policies and procedures related to cardiac monitoring.
- Provide appropriate ECG education for staff.
- Develop proficiency standards for all staff involved with ECG monitoring to ensure accurate and effective monitoring.

- Consider conducting an audit to assess:
  - Electrode placement
  - Lead selection

### **Need More Information or Help?**

1. Audit tool for measuring compliance with lead selection and lead placement available at [www.aacn.org](http://www.aacn.org)
2. Talk with a clinical practice specialist for additional information / assistance ([www.aacn.org](http://www.aacn.org), select PRN).

#### AACN Grading Level of Evidence

- Level I: Manufacturer's recommendations only
- Level II: Theory based, no research data to support recommendations:  
Recommendations from expert consensus group may exist
- Level III: Laboratory data, no clinical data to support recommendations
- Level IV: Limited clinical studies to support recommendations
- Level V: Clinical studies in more than one or two patient populations and situations to support recommendations
- Level VI: Clinical studies in a variety of patient populations and situations to support recommendations.

### **References:**

1. Thomason TR, Riegel B, Carlson B, Gocka I. Monitoring electrocardiographic changes: results of a national survey. *J Cardiovasc Nurs*. July 1995;9(4):1-9.
2. Drew BJ, Ide B, Sparacino PS. Accuracy of bedside electrocardiographic monitoring: a report on current practices of critical care nurses. *Heart Lung*. 1991;20(6):597-607.
3. Clochesy JM, Cifani L, Howe K. Electrode site preparation techniques: a follow-up study. *Heart Lung*. 1991;20(1):27-30.
4. Medina V, Clochesy JM, Omery A. Comparison of electrode site preparation techniques. *Heart Lung*. 1989;18(5):456-460.
5. Drew BJ. Celebrating the 100<sup>th</sup> birthday of the electrocardiogram: lessons learned from research in cardiac monitoring. *Am J Crit Care*. 2002;11(4):378-388.
6. Drew BJ, Ide B. Differential diagnosis of wide QRS complex tachycardia. *Prog Cardiovasc Nurs*. Summer 1998;13(3):46-47.
7. Drew BJ, Scheinman MM. ECG criteria to distinguish between aberrantly conducted supraventricular tachycardia and ventricular tachycardia: practical aspects for the immediate care setting. *Pacing Clin Electrophysiol*. 1995;18(12 pt 1):2194-2208.
8. Fabius DB. Diagnosing and treating ventricular tachycardia. *J Cardiovasc Nurs*. April 1993;7(3):8-25.
9. Drew BJ, Califf RM, Funk M, et al. Practice standards for electrocardiographic monitoring in hospital settings. *Circulation*. 2004;110(17):2721-2746. <http://circ.ahajournals.org/cgi/content/full/110/17/2721> Accessed April 21, 2008.
10. Drew BJ, Funk M. Practice standards for ECG monitoring in hospital settings: Executive summary and guide for implementation. *Crit Care Nurs Clin North Am*. 2006;18(2):157-168.
11. Drew BJ, Scheinman MM, Dracup K. MCL<sub>1</sub> and MCL<sub>6</sub> compared to V<sub>1</sub> and V<sub>6</sub> in distinguishing aberrant supraventricular from ventricular ectopic beats. *Pacing Clin Electrophysiol*. 1999;14(9):1375-1383.
12. Passman R, Kadish A. Polymorphic ventricular tachycardia, long Q-T syndrome, and torsades de pointes. *Med Clin North Am*. 2001;85(2):321-341.
13. Crouch MA, Limon L, Cassano AT. Clinical relevance and management of drug-related QT interval prolongation. *Pharmacotherapy*. 2003;23(7):881-908.
14. Sommargren CE, Drew BJ. Preventing torsades de pointes by careful cardiac monitoring in hospital settings. *AACN Adv Crit Care*. 2007;18(3):285-293.
15. Arizona Center for Education and Research on Therapeutics. [Drugs that Prolong the QT Interval and/or Induce Torsades de Pointes Ventricular Arrhythmia](http://www.qtdrugs.org/medical-pros/drug-lists/drug-lists.htm). 2006. <http://www.qtdrugs.org/medical-pros/drug-lists/drug-lists.htm> Accessed April 21, 2008.
16. Kern LS, McRae ME, Funk M. ECG monitoring after cardiac surgery: Postoperative atrial fibrillation and the atrial electrogram. *AACN Adv Crit Care*. 2007;18(3):294-304.
17. Miller JN, Drew BJ. Atrial electrograms after cardiac surgery: Survey of clinical practice. *Am J Crit Care*. 2007;16(4):350-356.
18. Mowery B, Suddaby EC. ECG Interpretation: what is different in children? *Pediatr Nurs*. 2001;27(3): 224-231.